

Review Article

PPARs: Key Regulators of Airway Inflammation and Potential Therapeutic Targets in Asthma

Asoka Banno¹, Aravind T. Reddy^{1,2}, Sowmya P. Lakshmi^{1,2}, and Raju C. Reddy^{1,2}

¹Department of Medicine, Division of Pulmonary, Allergy and Critical Care Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA 15213, USA

²Veterans Affairs Pittsburgh Healthcare System, Pittsburgh, PA 15240, USA

Abstract. Asthma affects approximately 300 million people worldwide, significantly impacting quality of life and healthcare costs. While current therapies are effective in controlling many patients' symptoms, a large number continue to experience exacerbations or treatment-related adverse effects. Alternative therapies are thus urgently needed. Accumulating evidence has shown that the peroxisome proliferator-activated receptor (PPAR) family of nuclear hormone receptors, comprising PPAR α , PPAR β/δ , and PPAR γ , is involved in asthma pathogenesis and that ligand-induced activation of these receptors suppresses asthma pathology. PPAR agonists exert their anti-inflammatory effects primarily by suppressing pro-inflammatory mediators and antagonizing the pro-inflammatory functions of various cell types relevant to asthma pathophysiology. Experimental findings strongly support the potential clinical benefits of PPAR agonists in the treatment of asthma. We review current literature, highlighting PPARs' key role in asthma pathogenesis and their agonists' therapeutic potential. With additional research and rigorous clinical studies, PPARs may become attractive therapeutic targets in this disease.

Keywords: PPAR, rosiglitazone, allergy, mucus, pulmonary

Corresponding Author

Raju C. Reddy
reddyrc@upmc.edu

Editor

Antigone Lazou

Dates

Received 4 August 2017
Accepted 11 December 2017

Copyright © 2018 Asoka Banno et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Asthma affects people of all ages worldwide, although its prevalence can vary widely depending on the specific demographics examined [1, 2]. An estimated 300 million individuals are affected by the disease, but its impact goes far beyond the patients themselves, involving families and communities and presenting a significant socioeconomic burden [1]. Clinically, asthma encompasses a heterogeneous group of phenotypes characterized by wheezing, coughing, dyspnea, chest tightness, and reduced expiratory airflow [2]. Pathologically, asthma is characterized by airway inflammation, remodeling, and hyperresponsiveness [3]. Underlying these anatomical and functional aberrations is the development of an abnormal T helper 2 (Th2) immune response [4, 5]. This response features an upsurge of Th2 lymphocytes that elevates production of interleukin-4 (IL-4), a cytokine promoting immunoglobulin E (IgE) synthesis, as well as of IL-5, which recruits eosinophils. Several other cell types and mediators are also involved in asthma pathogenesis. Airway epithelial cells, for instance, normally protect the lungs by serving as the first line of defense but, when impaired or dysregulated, contribute



to inflammation, remodeling, and mucus hypersecretion by producing vasoactive factors, pro-inflammatory agents, growth factors, and metalloproteinases. When the epithelium is compromised under pathological conditions, the interstitial tissue is also altered due to fibroblast proliferation and differentiation, collagen deposition, and hypertrophy and hyperplasia of airway smooth muscle cells that also produce pro-inflammatory factors [4, 5]. Another key player in asthma pathogenesis is the alveolar macrophage. As initial responders to external insults, these leukocytes, along with the airway epithelium, provide host defenses [4, 6] via their phagocytic function and secretion of appropriate molecules [7, 8]. Notably, to minimize subsequent tissue injury as well as to maintain healthy lung physiology and gas exchange, their regulation of immune responses is normally tightly controlled [4, 6, 9]. When dysregulation of their activities results in an imbalance between their anti- and pro-inflammatory responses [6, 8, 10], however, lung homeostasis is disrupted, as is seen in asthma. In fact, alterations of alveolar macrophage function have been observed in patients [4, 6].

Current standard therapies, most notably corticosteroids and β 2-adrenergic receptor agonists, effectively control symptoms and enhance lung function in many patients [11]. However, some individuals experience adverse events from these treatments while others face acute exacerbations without adequate improvement [5, 12]. These shortcomings of conventional treatments, combined with asthma's global burden, heighten the need for development of alternative, more effective therapies.

Peroxisome proliferator-activated receptors (PPARs), comprising PPAR α , PPAR β/δ , and PPAR γ , are nuclear hormone receptors initially recognized for their functions in lipid regulation and glucose metabolism [13]. As ligand-activated transcription factors ubiquitously expressed throughout the body [4, 14, 15], they are now known to also play a role in cellular processes such as differentiation, proliferation, survival, apoptosis, and motility in a variety of biological contexts including inflammation and immune responses [5, 16]. Cells of the immune system that infiltrate the airways following inflammatory stimuli (*e.g.* dendritic cells, eosinophils, macrophages, mast cells, monocytes, and neutrophils, as well as B and T lymphocytes) have been found to express PPARs [5]. Importantly, PPAR expression is altered during inflammatory responses, including airway inflammation, suggesting PPARs' involvement in asthma pathogenesis [4, 5]. Retrospective studies examining Chinese children [17] and adults [18] have provided further evidence by reporting correlations between certain PPAR single nucleotide polymorphisms and asthma risk and prognosis [17, 18]. These findings also highlight PPARs' potential as a predictive and prognostic molecular marker.

A variety of naturally occurring molecules and synthetic compounds activate PPARs. PPAR α agonists include polyunsaturated and saturated fatty acids and eicosanoids (*e.g.* 8(*S*)-hydroxyeicosatetraenoic acid and leukotriene B₄) as well as synthetic fibric acid derivatives (*e.g.* bezafibrate, clofibrate, and fenofibrate) and pirinixic acid (WY-14643) [5, 13, 16, 19]. Polyunsaturated and saturated fatty acids such as prostacyclin and other eicosanoids (*e.g.* prostaglandin A₁ and prostaglandin D₂) activate PPAR β/δ [5, 16]. Synthetic, high-affinity agonists for PPAR β/δ include GW501516, L165041, GW0742, and L783483 [4, 5, 20]. PPAR γ is stimulated by saturated and polyunsaturated fatty acids, eicosanoid derivatives such as 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ (15d-PGJ₂), and nitrated fatty acids [13, 14, 21, 22]. Thiazolidinediones (TZDs) such as pioglitazone, rosiglitazone, troglitazone, and ciglitazone are the most notable synthetic PPAR γ agonists [13].

Although studies have provided evidence for ligand-independent transcriptional activity, ligand-dependent functions of PPARs are better known and more widely accepted [23]. In this latter, conventional model, PPARs in their basal state are bound by corepressors that restrain their transcriptional activity [24]. PPAR agonists, however, trigger a conformational change that dissociates corepressors and favors coactivator interaction [25, 26]. The presence of coactivators accompanied by chromatin remodeling allows the receptors to heterodimerize with retinoid X receptors and bind to specific PPAR response elements (PPREs) in the promoters of their target genes, thus activating these genes' transcription [24–26]. Ligand binding also promotes ubiquitin-proteasome system-mediated degradation of corepressors [25].

In addition to corepressor/coactivator switches, post-translational modifications also regulate PPAR expression and activity. One such modification is phosphorylation, which can modulate PPARs' affinity for ligands, cofactors, retinoid X receptors, and target genes [23]. Depending on the cellular contexts and signals at play, phosphorylation can be stimulatory or suppressive [24]. Another post-translational modification is SUMOylation, which inhibits PPAR activity by promoting corepressor binding [24]. A third such modification is ubiquitination; ubiquitinated PPARs are subject to proteasomal degradation, thus downregulating their expression and activity [24].

Accumulating experimental evidence, with the majority focusing on PPAR α or PPAR γ , has shown that all three PPARs modulate the intensity, duration, and outcomes of inflammatory responses and that PPAR activation is anti-inflammatory and beneficial in various diseases associated with inflammation [4, 16, 27]. The cellular targets of this anti-inflammatory PPAR function are not only inflammatory cells of the immune system but also resident and structural cells of the airways that play significant roles during inflammation [4, 14].

At the molecular level, multiple mechanisms account for PPARs' anti-inflammatory effects. One such mechanism is coactivator sequestration: by competing for coactivators, PPARs limit the ability of pro-inflammatory transcription factors to access these required cofactors and initiate transcription of their target pro-inflammatory genes [13, 14]. PPARs can also inhibit inflammatory gene expression by stabilizing corepressor binding [24]. In addition, PPARs can directly bind to pro-inflammatory transcription factors, interfering with their access to coactivators or promoting corepressor recruitment, and consequently suppress their downstream gene transcription [24]. Transcription factors regulated this way by PPARs are major mediators of inflammatory responses and include activator protein-1 (AP-1), CCAAT/enhancer binding protein (C/EBP), nuclear factor of activated T cells (NFAT), nuclear factor- κ B (NF- κ B), and signal transducers and activators of transcription (STAT) [4]. Lastly, PPAR agonists have been shown to modulate c-Jun N-terminal kinase (JNK) and mitogen-activated protein kinase (MAPK) activities, indirectly suppressing inflammatory responses [24]. Besides these mechanisms, PPARs regulate expression of inflammatory modulators by binding to PPREs found in their promoters [28–30]. Thus, acting through pathways distinct from those employed by traditional therapies, PPAR-targeted asthma therapy could potentially prevent disease complications, progression, and exacerbations.

2. Roles of PPARs in Asthma

2.1. Overview

In general, expression and activity of each PPAR subtype is associated with protection against asthma or reduction in its severity, whereas impairment of a PPAR's function or expression leads to or exacerbates the disease. These effects target both inflammation and tissue remodeling, two prominent features of asthma. All three PPAR subtypes counteract inflammatory responses by modulating pro- and anti-inflammatory mediators as well as by reducing the expression of adhesion and chemotactic molecules essential for leukocyte recruitment. The specific molecules affected are not fully identical across the three subtypes, however.

PPARs also contribute to the preservation of tissue integrity in multiple ways. PPAR α and PPAR γ downregulate matrix metalloproteinases involved in extracellular matrix degradation, an essential aspect of tissue degradation and remodeling. PPAR β/δ and PPAR γ suppress lung fibroblasts' proliferation and their differentiation into myofibroblasts, further blocking increased collagen deposition. PPAR γ also inhibits epithelial and smooth muscle hyperplasia as well as blocking mucus overproduction. In the following sections, these multifaceted anti-asthma functions of each PPAR subtype are discussed in more detail.

2.2. PPAR α

PPAR α was first shown to control the duration of inflammatory responses in a mouse ear-swelling model [31]. *In vitro* and *in vivo* studies have since identified a variety of mechanisms by which PPAR α exerts its anti-inflammatory effect, including antagonism of inflammatory cell functions. For example, WY-14643 promotes apoptosis of human monocyte-derived macrophages [32]. PPAR α activation also reduces production of multiple pro-inflammatory mediators, including tumor necrosis factor- α (TNF- α), IL-1 α , IL-6, and IL-8, in multiple skin inflammation models [33, 34]. It also reduces production of TNF- α and IL-6 by monocytes *in vitro* [35], and of IL-6 and IL-8 by aortic smooth muscle cells in atherosclerosis models [36, 37]. In chronic inflammatory conditions, such as those characterized by constitutive NF- κ B activation and elevated levels of pro-inflammatory cytokines, WY-14643 treatment similarly suppresses TNF- α and IL-6 production [38]. Conversely, PPAR α deficiency exacerbates inflammatory features such as IL-6 and IL-12 production [39]. Furthermore, Ye *et al.* found that fenofibrate treatment reduced TNF- α and IL-6 levels in individuals with hypertriglyceridemia, a condition often associated with increased inflammatory markers [40].

In addition to suppressing pro-inflammatory cytokines, PPAR α activation controls expression or production of adhesion and chemotactic molecules that are imperative to inflammatory responses. Highlighting the essential role of PPAR α in migration, adhesion, and recruitment of immune cells, Michalik *et al.* showed that skin wound healing, where such migration is beneficial, was impaired in PPAR α -deficient mice [41]. Conversely, WY-14643 hinders pro-inflammatory neutrophil infiltration by suppressing intercellular adhesion molecule-1 (ICAM-1) expression in gingivomucosal tissues of rats with periodontitis [42], as well as in inflamed colons of mice with inflammatory bowel disease [43]. In the latter study, PPAR α knockout mice showed signs of more severe colonic injury than did wild-type animals. Vascular cell

adhesion molecule-1 (VCAM-1) expression in human aortic endothelial cells [44] and human carotid artery endothelial cells [45] is similarly reduced by WY-14643 and fenofibrate in *in vitro* inflammation models, consequently suppressing monocyte/macrophage binding to such cells. WY-14643 and fenofibrate treatments likewise reduce monocyte chemoattractant protein-1 (MCP-1) secretion from human umbilical vein endothelial cells [46].

Matrix metalloproteinases (MMPs), particularly MMP-9, contribute to inflammation by promoting extracellular matrix degradation during tissue remodeling associated with chronic inflammation and also by assisting infiltration of inflammatory cells through the basement membrane [4]. WY-14643 reduces MMP-9 expression in rat mesangial cells [47], while fenofibrate similarly decreases MMP-9 secretion by human monocytic cells [48].

In addition to inhibiting expression and activity of pro-inflammatory agents, activated PPAR α can induce anti-inflammatory agents. For example, fenofibrate increases IL-10 expression during experimental autoimmune myocarditis in mice [49] and WY-14643 promotes expression of anti-inflammatory sIL-1 receptor antagonist (sIL-1ra) [50]. Furthermore, WY-14643, fibrates, and another PPAR α agonist, GW9578, are known to induce I κ B α expression, thereby hindering NF- κ B's pro-inflammatory activity [51, 52]. Together these studies demonstrate that PPAR α controls inflammatory responses not only via downregulation of pro-inflammatory molecules but also via upregulation of anti-inflammatory mediators.

Consistent with the above findings in other organs and disease models, current data support the anti-inflammatory effect of PPAR α activation in the lungs and the airways. In murine models of allergic airway disease, PPAR α deficiency exacerbates asthmatic features such as airway hyperresponsiveness and eosinophilia, while treatment with PPAR α agonists shows the opposite trend [53–55]. In an experimental model of pleurisy, clofibrate treatment adds to the anti-inflammatory activity of the synthetic glucocorticoid dexamethasone [56]; combination therapy significantly downregulates macrophage and other inflammatory cell infiltration into the pleural cavity and thereby reduces tissue injury. Conversely, the absence of PPAR α compromises dexamethasone's control of lung inflammation in mice [56]. Thus, PPAR α agonists not only have the potential to be useful as monotherapy but also may function synergistically with glucocorticoids in asthma treatment. Together, these studies support the anti-inflammatory effects of PPAR α activation and justify further investigation of the receptor's role in asthma and airway inflammation.

2.3. PPAR β/δ

Studies using high-affinity ligands such as GW501516 and GW0742 have shown that PPAR β/δ modulates many mediators of inflammation [57–60]. In monocytes/macrophages, this anti-inflammatory function of PPAR β/δ rests in part on ligand binding-induced dissociation from the transcriptional repressor B cell lymphoma-6 (Bcl-6) protein; this uncoupling releases Bcl-6 to suppress expression of pro-inflammatory molecules [57, 60]. More directly, PPAR β/δ activation by GW501516 antagonizes inflammation by inducing expression of sIL-1ra [61] and transforming growth factor- β 1 (TGF- β 1) [62]. Like PPAR α - and PPAR γ -activating ligands, PPAR β/δ agonists suppress endothelial cells' expression of adhesion molecules such as VCAM-

1, ICAM-1, and E-selectin that are required for leukocyte recruitment [63–66] as well as the chemokines MCP-1 and growth-regulated oncogene- α (GRO α) [63, 65, 66].

PPAR β/δ is also involved in wound healing-relevant functions of keratinocytes, which express PPAR β/δ more abundantly than the other PPAR isotypes [67]. PPAR β/δ upregulates anti-apoptotic genes and downregulates pro-apoptotic genes, resulting in keratinocyte survival [68]. The activated receptor further enhances wound healing both by potentiating keratinocytes' migratory response to injury via enhancement of chemotactic signals and by promoting integrin recycling and actin cytoskeleton remodeling [69]. An *in vivo* study has validated this conclusion by showing that PPAR β/δ -deficient mice exhibit an impaired wound-healing response [41].

A PPAR β/δ ligand was initially shown ineffective in controlling allergen-induced airway inflammation in mice [55]. However, a later study demonstrated that GW0742 inhibits lipopolysaccharide-induced neutrophil infiltration into lung tissues and hinders production of IL-6, IL-1 β , and TNF α , thus diminishing the extent of inflammatory responses [70]. Furthermore, GW0742 blocks pulmonary fibroblast proliferation [71] and controls leukocyte infiltration and tissue damage in a mouse model of pulmonary fibrosis [72]; subepithelial fibrosis is a prominent component of airway remodeling during asthma pathogenesis. Of note, the discrepancy in findings between Trifilieff *et al.* and Haskova *et al.* may result from differences in the timing of PPAR β/δ agonist administration. Alternatively, the observed disagreement may reflect use of different disease models. In summary, although accumulating evidence supports the anti-inflammatory properties of PPAR β/δ agonists, additional studies are needed to elucidate the role of PPAR β/δ in airway inflammation and to assess its prospect as a therapeutic target for asthma.

2.4. PPAR γ

PPAR γ 's expression by various cells of the immune system underscores its prominent role in inflammatory responses [14, 16, 73, 74]. Following initial recognition as a regulator of monocytes/macrophage function in atherosclerosis [73], PPAR γ is now known to regulate functions of other inflammation-associated cell types [4, 13, 74, 75] in various disease and disease model contexts [4, 73, 75]. Furthermore, many inflammatory conditions are associated with alterations in PPAR γ expression and activity, and such changes are believed to contribute significantly to several diseases [5, 76]. As its involvement in inflammation has been extensively reviewed elsewhere [14, 16, 73], the focus in this review will be placed on PPAR γ 's role in asthma.

IL-4, a cytokine that promotes the Th2 responses associated with asthma pathogenesis, induces PPAR γ in airway epithelial cells [77]. To substantiate this *in vitro* finding, studies using a murine model of allergic airway disease observed higher levels of PPAR γ in the lung tissues of animals exposed to the allergen ovalbumin (OVA) [78–80]. This upregulation of PPAR γ was localized to airway epithelial cells, smooth muscle cells, mast cells, and some inflammatory cells [80]. The link between asthma pathogenesis and PPAR γ expression levels is emphasized by a study showing that asthmatic patients exhibit greater PPAR γ expression in their bronchial submucosa, bronchial epithelium, and airway smooth muscle than do healthy controls, and that this upregulation is reversed by glucocorticoid treatment [81]. It has been speculated that increased PPAR γ expression is a cellular response to pro-inflammatory cytokines that initiates a negative feedback pathway limiting airway inflammation [5]. In contrast, alveolar macrophages

of allergen-challenged asthmatic patients were shown to have reduced PPAR γ levels compared to those in controls [82]. The authors suggest that this downregulation could potentially contribute to airway inflammation. Alternatively, the findings by Honda *et al.* showing that the increase in PPAR γ expression in allergen-sensitized and -challenged animals was blocked by ciglitazone treatment [80] offer another plausible explanation: this PPAR γ downregulation from otherwise elevated levels may be a consequence of PPAR γ activation-induced reduction or resolution of airway inflammation [4, 5]. Thus, while PPAR γ levels appear to influence asthma pathogenesis, analysis and interpretation of expression data must include careful consideration of the complex interaction between PPAR γ and the stage of inflammation (*i.e.* initiation vs. resolution) [5].

PPAR γ activation/PPAR γ agonists have displayed beneficial effects on multiple asthma features. For example, in a mouse model of OVA-induced allergic airway disease, rosiglitazone reduced airway hyperresponsiveness [83]. In another mouse model, which induces allergic airway disease via cockroach allergen, pioglitazone demonstrated the same effect as well as suppression of leukocyte infiltration, pro-inflammatory chemokine and cytokine production, and mucus overproduction [84]. Importantly, effects on pathophysiological responses and cytokine and chemokine production were comparable between pioglitazone and dexamethasone. Furthermore, ciglitazone significantly suppresses airway inflammation and remodeling in addition to airway hyperresponsiveness, eosinophilia, mucus overproduction, cytokine production, and collagen deposition [53, 80, 85]. Yet another PPAR γ agonist, troglitazone, inhibits IL-5-mediated survival and eotaxin-directed chemotaxis of eosinophils [86], indicating its efficacy against eosinophilia. Significantly, Mueller *et al.* reported that ciglitazone administered later in the course of allergen exposure is also effective in reducing airway inflammation, as suggested by decrease in inflammatory cell infiltration and epithelial hyperplasia in the lungs [85].

PPAR γ agonists suppress functions of inflammatory cells other than eosinophils. Rosiglitazone decreases lymph node infiltration of lung dendritic cells, critical inducers of immune responses, in OVA-treated animals [87, 88], and thus reduces airway inflammation [88]. OVA-induced inflammation assessed by bronchoalveolar lavage is also reduced by the synthetic PPAR γ agonist GI262570 [55]. In addition to suppressing pro-inflammatory cytokine production, 15d-PGJ₂ and troglitazone enhanced phagocytosis of apoptotic neutrophils by human alveolar macrophages, an important aspect of inflammatory resolution [89]. These macrophages also show upregulated CD36 expression after PPAR γ agonist treatment. Consistently, another study using a bleomycin-induced lung fibrosis model reported that enhancement of alveolar macrophages' efferocytotic ability in the presence of apoptotic cells was reversed by the PPAR γ antagonist GW9662 [90], emphasizing the prominent role of PPAR γ in macrophage regulation.

PPAR γ activation also regulates structural cells involved in airway inflammation. An *in vitro* study showed both 15d-PGJ₂ and ciglitazone inhibited proliferation and induced apoptosis of human airway smooth muscle cells, whose hypertrophy and hyperplasia contribute significantly to asthma-associated airway narrowing [91]. In addition, rosiglitazone and pioglitazone have been shown to reduce MMP-9 activity and protein expression in TNF- α - or phorbol 12-myristate 13-acetate (PMA)-stimulated human bronchial epithelial cells [92], thus suggesting their efficacy against the tissue remodeling observed during asthma pathogenesis.

Airway smooth muscle cells also contribute to inflammation by secreting granulocyte-macrophage colony-stimulating factor (GM-CSF), a cytokine critical for survival and activity

of various leukocytes, including eosinophils [91], and 15d-PGJ₂ and ciglitazone suppress GM-CSF release [91]. This finding provides further evidence for the effectiveness of PPAR γ agonists against inflammation. Moreover, 15d-PGJ₂ and ciglitazone downregulate IL-8 secretion from airway epithelial cells [77], which is expected to primarily reduce neutrophil recruitment during airway inflammation. PPAR γ agonists also decrease lung expression of ICAM-1 and VCAM-1 as well as levels of eotaxin and regulated upon activation normal T cell expressed and secreted (RANTES) in a mouse model of occupational asthma [93].

Lung injury, especially to the alveolar epithelium, induces fibroblasts to proliferate and differentiate into myofibroblasts that produce excessive collagen and other extracellular matrix components [4, 94–96]. 15d-PGJ₂, troglitazone, ciglitazone, and rosiglitazone [95, 96] as well as constitutively active PPAR γ [95] prevent human lung fibroblasts from differentiating into myofibroblasts [95, 96]. PPAR γ agonists also suppress collagen secretion from these cells [95, 96] and inhibit bleomycin-induced pulmonary fibrosis [95]. Taken together, the studies cited in this section provide evidence for multifaceted anti-inflammatory effects of PPAR γ in the lungs and suggest that its agonists may become useful in asthma intervention.

3. Therapeutic Implication of PPAR Ligands for Asthma

Collectively, experimental findings strongly support the clinical benefits of PPAR agonists as asthma treatments. Unfortunately, however, clinical data are currently available only for PPAR γ agonists. Supporting the value of PPAR γ agonists as asthma therapy, a recent retrospective cohort study analyzing a large number of diabetic patients with asthma found an association between TZD use (for diabetes treatment) and reduction in the risk of asthma exacerbations as well as in oral steroid prescriptions [97]. Another study of 16 steroid-naïve asthmatic patients also reported that 12 weeks of rosiglitazone treatment improved airway hyperresponsiveness (assessed by response to methacholine) [98]. In agreement with these clinical studies, a case report described improvement of asthma with pioglitazone treatment [99]: a 71-year-old man with type 2 diabetes, hyperlipidemia, hypertension, and asthma experienced disappearance of wheezing after several days of pioglitazone treatment. Another diabetic man with asthma also showed similar clinical improvement [99]. Moreover, upon discontinuation of pioglitazone, his respiratory symptoms returned, emphasizing the association between pioglitazone treatment and recovery from asthma.

Still, the efficacy of PPAR γ agonists as asthma drugs remains controversial. A randomized study of 46 asthmatic patients failed to observe improvement in asthma symptoms (assessed by Asthma Control Questionnaire score) after 4-week rosiglitazone treatment, although it did find that the treatment enhanced patients' lung function [100]. Likewise, in a placebo-controlled, randomized study of 32 asthma patients, 4-week rosiglitazone treatment only modestly decreased late phase asthma reactivity to allergen challenge, leading the authors to conclude that rosiglitazone would not provide adequate intervention [101]. A double-blind, randomized controlled trial of 68 asthma patients reached a similar conclusion after observing no sign of improvement after 12 weeks of pioglitazone [102]. It is noteworthy, however, that all these studies are associated with some limitations such as a small sample size and non-general subjects. Thus, larger randomized, placebo-controlled studies should be conducted with various types of asthma patients to substantiate the clinical effects of PPAR γ agonists. Similar studies

on the use of PPAR α and PPAR β/δ agonists can be expected to provide further insights into asthma treatment.

4. Conclusions

Traditional asthma therapies, although effective for many patients, provide only temporary symptomatic alleviation [103]. Moreover, even with these interventions, some patients still experience exacerbations and progressive deterioration of pulmonary function [5, 12]. Understanding the fundamental pathophysiology is thus critical for advances in asthma therapy. Accumulating experimental findings support all three PPARs' anti-inflammatory properties and involvement in airway inflammation. Clinical data available for PPAR γ agonists also substantiate their therapeutic potential. Unfortunately, PPAR γ agonists are associated with side effects: rosiglitazone and pioglitazone have been shown to cause weight gain, edema, and congestive heart failure [104] as well as bone fractures [105]; troglitazone is associated with hepatotoxicity and has therefore been withdrawn from clinical use [106]. Thus, to minimize these adverse effects, the drugs may be best administered via inhalation as opposed to systemic delivery [27]. Importantly, local administration of the drugs in murine models has been shown to provide similar benefits to those seen with systemic delivery on multiple pathological features of asthma, including elevated cytokine production, airway hyperresponsiveness, and eosinophilia, [53, 55], thus supporting inhalational drug delivery.

Another strategy to circumvent or reduce the side effects associated with PPAR agonists or traditional therapies is to employ combinations of drugs, each at a lower concentration that may offer limited benefit as monotherapy. In fact, PPAR γ agonists have demonstrated synergistic effects with corticosteroids and β 2-adrenergic receptor agonists. In a mouse model of inflammation, individually ineffective doses of rosiglitazone and dexamethasone reduced paw edema when administered together [107]. The β 2-adrenergic receptor agonist salbutamol also displayed synergy with 15d-PGJ₂ or rosiglitazone in reduction of human bronchial smooth muscle cell proliferation [108]. Additive inhibition of TNF- α -induced chemokine production was similarly observed with 15d-PGJ₂ and the glucocorticoid fluticasone as well as 15d-PGJ₂ and the β 2-adrenergic receptor agonist salmeterol [109]. As noted, clofibrate also showed synergy with dexamethasone in a mouse pleurisy model [56]. These data suggest combination therapy may be an attractive option. Thus, with further investigation and clinical trials, PPAR agonists may become an effective part of asthma therapy.

Abbreviations

15d-PGJ₂: 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂

AP-1: Activator protein-1

Bcl-6: B-cell lymphoma-6

C/EBP: CCAAT/enhancer binding protein

GM-CSF: Granulocyte-macrophage colony-stimulating factor

GRO- α : Growth-regulated oncogene- α

ICAM-1: Intercellular adhesion molecule-1

IgE: Immunoglobulin E
IL: Interleukin
JNK: c-Jun N-terminal kinase
MAPK: mitogen-activated protein kinase
MCP-1: Monocyte chemotactic protein-1
MMP: Matrix metalloproteinase
NFAT: Nuclear factor of activated T cells
NF- κ B: Nuclear factor- κ B
OVA: Ovalbumin
PMA: Phorbol 12-myristate 13-acetate
PPAR: Peroxisome proliferator-activated receptor
PPRE: PPAR response element
RANTES: Regulated upon activation normal T cell expressed and secreted
sIL-1ra: Secreted IL-1 receptor antagonist
STAT: Signal transducers and activators of transcription
TGF- β 1: Transforming growth factor- β 1
Th2: T helper 2
TNF- α : Tumor necrosis factor- α
TZD: Thiazolidinedione
VCAM-1: Vascular cell adhesion molecule-1

Competing Interests

The authors declare no competing interests.

Acknowledgments

This work was supported by a Merit Review award from the U.S. Department of Veterans Affairs and National Institutes of Health grants HL093196 and AI125338 (RCR).

Disclaimer

The contents in this article do not represent the views of the U.S. Department of Veterans Affairs or the United States Government.

References

- [1] T. To, S. Stanojevic, G. Moores et al., "Global asthma prevalence in adults: findings from the cross-sectional world health survey," *BMC Public Health*, vol. 12, article 204, 2012.
- [2] H. K. Reddel, E. D. Bateman, A. Becker et al., "A summary of the new GINA strategy: a roadmap to asthma control," *European Respiratory Journal*, vol. 46, no. 3, pp. 622–639, 2015.

- [3] C. Donovan, X. Tan, and J. E. Bourke, "PPAR γ ligands regulate noncontractile and contractile functions of airway smooth muscle: Implications for asthma therapy," *PPAR Research*, Article ID 809164, 2012.
- [4] J. Becker, C. Delayre-Orthez, N. Frossard, and F. Pons, "Regulation of inflammation by PPARs: A future approach to treat lung inflammatory diseases?" *Fundamental & Clinical Pharmacology*, vol. 20, no. 5, pp. 429–447, 2006.
- [5] J. E. Ward and X. Tan, "Peroxisome proliferator activated receptor ligands as regulators of airway inflammation and remodelling in chronic lung disease," *PPAR Research*, vol. 2007, Article ID 14983, 12 pages, 2007.
- [6] J. Balhara and A. S. Gounni, "The alveolar macrophages in asthma: A double-edged sword," *Mucosal Immunology*, vol. 5, no. 6, pp. 605–609, 2012.
- [7] K. Hiraiwa and S. F. van Eeden, "Contribution of lung macrophages to the inflammatory responses induced by exposure to air pollutants," *Mediators of Inflammation*, vol. 2013, Article ID 619523, 10 pages, 2013.
- [8] J. B. Rubins, "Alveolar macrophages: Wielding the double-edged sword of inflammation," *American Journal of Respiratory and Critical Care Medicine*, vol. 167, no. 2, pp. 103–104, 2003.
- [9] J. D. Aberdein, J. Cole, M. A. Bewley, H. M. Marriott, and D. H. Dockrell, "Alveolar macrophages in pulmonary host defence—the unrecognized role of apoptosis as a mechanism of intracellular bacterial killing," *Clinical & Experimental Immunology*, vol. 174, no. 2, pp. 193–202, 2013.
- [10] M. Peters-Golden, "The alveolar macrophage: The forgotten cell in asthma," *American Journal of Respiratory Cell and Molecular Biology*, vol. 31, no. 1, pp. 3–7, 2004.
- [11] J. Agbetile and R. Green, "New therapies and management strategies in the treatment of asthma: Patient-focused developments," *Journal of Asthma and Allergy*, no. 4, pp. 1–12, 2011.
- [12] J. O'Toole, L. Mikulic, and D. A. Kaminsky, "Epidemiology and Pulmonary Physiology of Severe Asthma," *Immunology and Allergy Clinics of North America*, vol. 36, no. 3, pp. 425–438, 2016.
- [13] M. G. Belvisi and J. A. Mitchell, "Targeting PPAR receptors in the airway for the treatment of inflammatory lung disease," *British Journal of Pharmacology*, vol. 158, no. 4, pp. 994–1003, 2009.
- [14] W. Wahli and L. Michalik, "PPARs at the crossroads of lipid signaling and inflammation," *Trends in Endocrinology & Metabolism*, vol. 23, no. 7, pp. 351–363, 2012.
- [15] S. Tyagi, P. Gupta, A. S. Saini, C. Kaushal, and S. Sharma, "The peroxisome proliferator-activated receptor: a family of nuclear receptors role in various diseases," *Journal of Advanced Pharmaceutical Technology & Research*, vol. 2, no. 4, pp. 236–240, 2011.
- [16] R. A. Daynes and D. C. Jones, "Emerging roles of PPARs in inflammation and immunity," *Nature Reviews Immunology*, vol. 2, no. 10, pp. 748–759, 2002.
- [17] Y. Zhang, Z. Wang, and T. Ma, "Associations of Genetic Polymorphisms Relevant to Metabolic Pathway of Vitamin D3 with Development and Prognosis of Childhood Bronchial Asthma," *DNA and Cell Biology*, vol. 36, no. 8, pp. 682–692, 2017.
- [18] W. Li, W. Dai, J. Sun et al., "Association of peroxisome proliferator-activated receptor- γ gene polymorphisms and gene-gene interaction with asthma risk in a Chinese adults population," *International Journal of Clinical and Experimental Medicine*, vol. 8, no. 10, pp. 19346–19352, 2015.
- [19] V. R. Narala, R. K. Adapala, M. V. Suresh, T. G. Brock, M. Peters-Golden, and R. C. Reddy, "Leukotriene B4 is a physiologically relevant endogenous peroxisome proliferator-activated receptor- α agonist," *The Journal of Biological Chemistry*, vol. 285, no. 29, pp. 22067–22074, 2010.
- [20] V. G. Keshamouni, S. Han, and J. Roman, "Peroxisome proliferator-activated receptors in lung cancer," *PPAR Research*, vol. 2007, Article ID 90289, 10 pages, 2007.
- [21] Y. Li, J. Zhang, and F. J. Schopfer, "Molecular recognition of nitrated fatty acids by PPAR gamma," *Nature Structural & Molecular Biology*, vol. 15, no. 8, pp. 865–867, 2008.
- [22] A. T. Reddy, S. P. Lakshmi, Y. Zhang, and R. C. Reddy, "Nitrated fatty acids reverse pulmonary fibrosis by dedifferentiating myofibroblasts and promoting collagen uptake by alveolar macrophages," *The FASEB Journal*, vol. 28, no. 12, pp. 5299–5310, 2014.
- [23] J. N. Feige, L. Gelman, L. Michalik, B. Desvergne, and W. Wahli, "From molecular action to physiological outputs: peroxisome proliferator-activated receptors are nuclear receptors at the crossroads of key cellular functions," *Progress in Lipid Research*, vol. 45, no. 2, pp. 120–159, 2006.
- [24] G. S. Harmon, M. T. Lam, and C. K. Glass, "PPARs and lipid ligands in inflammation and metabolism," *Chemical Reviews*, vol. 111, no. 10, pp. 6321–6340, 2011.
- [25] V. Perissi, A. Aggarwal, C. K. Glass, D. W. Rose, and M. G. Rosenfeld, "A corepressor/coactivator exchange complex required for transcriptional activation by nuclear receptors and other regulated transcription factors," *Cell*, vol. 116, no. 4, pp. 511–526, 2004.
- [26] J. N. Feige and J. Auwerx, "Transcriptional coregulators in the control of energy homeostasis," *Trends in Cell Biology*, vol. 17, no. 6, pp. 292–301, 2007.
- [27] R. C. Reddy, V. K. Rehan, J. Roman, and P. J. Sime, "PPARs: Regulators and translational targets in the lung," *PPAR Research*, Article ID 342924, 2012.

- [28] P. W. Thompson, A. I. Bayliffe, A. P. Warren, and J. R. Lamb, "Interleukin-10 is upregulated by nanomolar rosiglitazone treatment of mature dendritic cells and human CD4+ T cells," *Cytokine*, vol. 39, no. 3, pp. 184–191, 2007.
- [29] M. Zenhom, A. Hyder, I. Kraus-Stojanowic, A. Auinger, T. Roeder, and J. Schrezenmeir, "PPAR γ -dependent peptidoglycan recognition protein 3 (PGlyRP3) expression regulates proinflammatory cytokines by microbial and dietary fatty acids," *Immunobiology*, vol. 216, no. 6, pp. 715–724, 2011.
- [30] D. A. Mogilenko, I. V. Kudriavtsev, V. S. Shavva et al., "Peroxisome proliferator-activated receptor α positively regulates complement C3 expression but inhibits tumor necrosis factor-mediated activation of C3 gene in mammalian hepatic-derived cells," *The Journal of Biological Chemistry*, vol. 288, no. 3, pp. 1726–1738, 2013.
- [31] P. R. Devchand, H. Keller, J. M. Peters, M. Vazquez, F. J. Gonzalez, and W. Wahli, "The PPAR α -leukotriene B4 pathway to inflammation control," *Nature*, vol. 384, no. 6604, pp. 39–43, 1996.
- [32] G. Chinetti, S. Griglio, M. Antonucci et al., "Activation of proliferator-activated receptors α and γ induces apoptosis of human monocyte-derived macrophages," *The Journal of Biological Chemistry*, vol. 273, no. 40, pp. 25573–25580, 1998.
- [33] M. Y. Sheu, A. J. Fowler, J. Kao et al., "Topical peroxisome proliferator activated receptor- α activators reduce inflammation in irritant and allergic contact dermatitis models," *Journal of Investigative Dermatology*, vol. 118, no. 1, pp. 94–101, 2002.
- [34] S. Kippenberger, S. M. Loitsch, M. Grundmann-Kollmann et al., "Activators of peroxisome proliferator-activated receptors protect human skin from ultraviolet-B-light-induced inflammation," *Journal of Investigative Dermatology*, vol. 117, no. 6, pp. 1430–1436, 2001.
- [35] C. K. Combs, P. Bates, J. C. Karlo, and G. E. Landreth, "Regulation of beta-amyloid stimulated proinflammatory responses by peroxisome proliferator-activated receptor alpha," *Neurochemistry International*, vol. 39, no. 5-6, pp. 449–457, 2001.
- [36] B. Staels, W. Koenig, A. Habib et al., "Activation of human aortic smooth-muscle cells is inhibited by PPAR α but not by PPAR γ activators," *Nature*, vol. 393, no. 6687, pp. 790–793, 1998.
- [37] S. Ryoo, M. Won, D.-U. Kim et al., "PPAR α activation abolishes LDL-stimulated IL-8 production via AP-1 deactivation in human aortic smooth muscle cells," *Biochemical and Biophysical Research Communications*, vol. 318, no. 2, pp. 329–334, 2004.
- [38] N. F. L. Spencer, M. E. Poynter, S.-Y. Im, and R. A. Daynes, "Constitutive activation of NF- κ B in an animal model of aging," *International Immunology*, vol. 9, no. 10, pp. 1581–1588, 1997.
- [39] M. E. Poynter and R. A. Daynes, "Peroxisome proliferator-activated receptor α activation modulates cellular redox status, represses nuclear factor- κ B signaling, and reduces inflammatory cytokine production in aging," *The Journal of Biological Chemistry*, vol. 273, no. 49, pp. 32833–32841, 1998.
- [40] P. Ye, J.-J. Li, G. Su, and C. Zhang, "Effects of fenofibrate on inflammatory cytokines and blood pressure in patients with hypertriglyceridemia [3]," *Clinica Chimica Acta*, vol. 356, no. 1-2, pp. 229–232, 2005.
- [41] L. Michalik, B. Desvergne, N. S. Tan et al., "Impaired skin wound healing in peroxisome proliferator-activated receptor (PPAR) α and PPAR β mutant mice," *The Journal of Cell Biology*, vol. 154, no. 4, pp. 799–814, 2001.
- [42] E. Briguglio, R. Di Paola, I. Paterniti et al., "WY-14643, a potent peroxisome proliferator activator receptor- α PPAR- α agonist ameliorates the inflammatory process associated to experimental periodontitis," *PPAR Research*, Article ID 193019, 2010.
- [43] S. Cuzzocrea, R. Di Paola, E. Mazzon et al., "Role of endogenous and exogenous ligands for the peroxisome proliferators activated receptors alpha (PPAR- α) in the development of inflammatory bowel disease in mice," *Laboratory Investigation*, vol. 84, no. 12, pp. 1643–1654, 2004.
- [44] S. M. Jackson, F. Parhami, X.-P. Xi et al., "Peroxisome proliferator-activated receptor activators target human endothelial cells to inhibit leukocyte-endothelial cell interaction," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 19, no. 9, pp. 2094–2104, 1999.
- [45] N. Marx, G. K. Sukhova, T. Collins, P. Libby, and J. Plutzky, "PPAR α activators inhibit cytokine-induced vascular cell adhesion molecule-1 expression in human endothelial cells," *Circulation*, vol. 99, no. 24, pp. 3125–3131, 1999.
- [46] V. Pasceri, J. Chang, J. T. Willerson, and E. T. H. Yeh, "Modulation of C-reactive protein-mediated monocyte chemoattractant protein-1 induction in human endothelial cells by anti-atherosclerosis drugs," *Circulation*, vol. 103, no. 21, pp. 2531–2534, 2001.
- [47] W. Eberhardt, E. L. S. Akool, J. Rebhan et al., "Inhibition of cytokine-induced matrix metalloproteinase 9 expression by peroxisome proliferator-activated receptor α agonists is indirect and due to a NO-mediated reduction of mRNA stability," *The Journal of Biological Chemistry*, vol. 277, no. 36, pp. 33518–33528, 2002.
- [48] H. Shu, B. Wong, G. Zhou et al., "Activation of PPAR α or γ reduces secretion of matrix metalloproteinase 9 but not interleukin 8 from human monocytic THP-1 cells," *Biochemical and Biophysical Research Communications*, vol. 267, no. 1, pp. 345–349, 2000.
- [49] S. Maruyama, K. Kato, M. Kodama et al., "Fenofibrate, a peroxisome proliferator-activated receptor alpha activator, suppresses experimental autoimmune myocarditis by stimulating the interleukin-10 pathway in rats," *Journal of Atherosclerosis and Thrombosis*, vol. 9, no. 2, pp. 87–92, 2002.

- [50] R. Stienstra, S. Mandard, N. S. Tan et al., "The Interleukin-1 receptor antagonist is a direct target gene of PPAR α in liver," *Journal of Hepatology*, vol. 46, no. 5, pp. 869–877, 2007.
- [51] P. Delerive, P. Gervois, J.-C. Fruchart, and B. Staels, "Induction of I κ B α expression as a mechanism contributing to the anti-inflammatory activities of peroxisome proliferator-activated receptor- α activators," *The Journal of Biological Chemistry*, vol. 275, no. 47, pp. 36703–36707, 2000.
- [52] P. Delerive, K. De Bosscher, W. V. Berghe, J.-C. Fruchart, G. Haegeman, and B. Staels, "DNA binding-independent induction of I κ B α gene transcription by PPAR α ," *Molecular Endocrinology*, vol. 16, no. 5, pp. 1029–1039, 2002.
- [53] G. Woerly, K. Honda, M. Loyens et al., "Peroxisome proliferator-activated receptors α and γ down-regulate allergic inflammation and eosinophil activation," *The Journal of Experimental Medicine*, vol. 198, no. 3, pp. 411–421, 2003.
- [54] C. Delayre-Orthez, J. Becker, J. Auwerx, N. Frossard, and F. Pons, "Suppression of allergen-induced airway inflammation and immune response by the peroxisome proliferator-activated receptor- α agonist fenofibrate," *European Journal of Pharmacology*, vol. 581, no. 1-2, pp. 177–184, 2008.
- [55] A. Trifilieff, A. Bench, M. Hanley, D. Bayley, E. Campbell, and P. Whittaker, "PPAR- α and - γ but not - δ agonists inhibit airway inflammation in a murine model of asthma: In vitro evidence for an NF- κ B-independent effect," *British Journal of Pharmacology*, vol. 139, no. 1, pp. 163–171, 2003.
- [56] S. Cuzzocrea, S. Bruscoli, E. Mazzon et al., "Peroxisome proliferator-activated receptor- α contributes to the anti-inflammatory activity of glucocorticoids," *Molecular Pharmacology*, vol. 73, no. 2, pp. 323–337, 2008.
- [57] C. Lee, A. Chawla, N. Urbiztondo, D. Liao, W. A. Boisvert, and R. M. Evans, "Transcriptional repression of atherogenic inflammation: modulation by PPAR δ ," *Science*, vol. 302, no. 5644, pp. 453–457, 2003.
- [58] A. C. Li, C. J. Binder, A. Gutierrez et al., "Differential inhibition of macrophage foam-cell formation and atherosclerosis in mice by PPAR α , β/δ , and γ ," *The Journal of Clinical Investigation*, vol. 114, no. 11, pp. 1564–1576, 2004.
- [59] G. D. Barish, A. R. Atkins, and M. Downes, "PPAR δ regulates multiple proinflammatory pathways to suppress atherosclerosis," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 11, pp. 4271–4276, 2008.
- [60] Y. Takata, J. Liu, F. Yin et al., "PPAR δ -mediated antiinflammatory mechanisms inhibit angiotensin II-accelerated atherosclerosis," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 11, pp. 4277–4282, 2008.
- [61] H. C. Chong, M. J. Tan, V. Philippe et al., "Regulation of epithelial-mesenchymal IL-1 signaling by PPAR β/δ is essential for skin homeostasis and wound healing," *The Journal of Cell Biology*, vol. 184, no. 6, pp. 817–831, 2009.
- [62] H. J. Kim, S. A. Ham, S. U. Kim et al., "Transforming growth factor- β 1 is a molecular target for the peroxisome proliferator-activated receptor δ ," *Circulation Research*, vol. 102, no. 2, pp. 193–200, 2008.
- [63] Y. Rival, N. Benéteau, T. Taillandier et al., "PPAR α and PPAR δ activators inhibit cytokine-induced nuclear translocation of NF- κ B and expression of VCAM-1 in EAhy926 endothelial cells," *European Journal of Pharmacology*, vol. 435, no. 2-3, pp. 143–151, 2002.
- [64] Y. Fan, Y. Wang, Z. Tang et al., "Suppression of pro-inflammatory adhesion molecules by PPAR- δ in human vascular endothelial cells," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 28, no. 2, pp. 315–321, 2008.
- [65] L. Piqueras, M. J. Sanz, M. Perretti et al., "Activation of PPAR β/δ inhibits leukocyte recruitment, cell adhesion molecule expression, and chemokine release," *Journal of Leukocyte Biology*, vol. 86, no. 1, pp. 115–122, 2009.
- [66] Y.-J. Liang, Y.-C. Liu, C.-Y. Chen et al., "Comparison of PPAR δ and PPAR γ in inhibiting the pro-inflammatory effects of C-reactive protein in endothelial cells," *International Journal of Cardiology*, vol. 143, no. 3, pp. 361–367, 2010.
- [67] M. Westergaard, J. Henningsen, M. L. Svendsen et al., "Modulation of keratinocyte gene expression and differentiation by PPAR-selective ligands and tetradecylthioacetic acid," *Journal of Investigative Dermatology*, vol. 116, no. 5, pp. 702–712, 2001.
- [68] N. S. Tan, L. Michalik, N. Noy et al., "Critical roles of PPARbeta/delta in keratinocyte response to inflammation," *Genes & Development*, vol. 15, no. 24, pp. 3263–3277, 2001.
- [69] N. S. Tan, G. Icre, A. Montagner, B. Bordier-ten-Heggeler, W. Wahli, and L. Michalik, "The nuclear hormone receptor peroxisome proliferator-activated receptor β/δ potentiates cell chemotaxis, polarization, and migration," *Molecular and Cellular Biology*, vol. 27, no. 20, pp. 7161–7175, 2007.
- [70] Z. Haskova, B. Hoang, G. Luo et al., "Modulation of LPS-induced pulmonary neutrophil infiltration and cytokine production by the selective PPAR β/δ ligand GW0742," *Inflammation Research*, vol. 57, no. 7, pp. 314–321, 2008.
- [71] F. Y. Ali, K. Egan, G. A. FitzGerald et al., "Role of prostacyclin versus peroxisome proliferator-activated receptor beta receptors in prostacyclin sensing by lung fibroblasts," *American Journal of Respiratory Cell and Molecular Biology*, vol. 34, no. 2, pp. 242–246, 2006.

- [72] M. Galuppo, R. di Paola, E. Mazzon et al., "GW0742, a high affinity PPAR- β/δ agonist reduces lung inflammation induced by bleomycin instillation in mice," *International Journal of Immunopathology and Pharmacology*, vol. 23, no. 4, pp. 1033–1046, 2010.
- [73] R. B. Clark, "The role of PPARs in inflammation and immunity," *Journal of Leukocyte Biology*, vol. 71, no. 3, pp. 388–400, 2002.
- [74] A. Croasdell, P. F. Duffney, N. Kim, S. H. Lacy, P. J. Sime, and R. P. Phipps, "PPAR γ and the innate immune system mediate the resolution of inflammation," *PPAR Research*, vol. 2015, Article ID 549691, 20 pages, 2015.
- [75] M. V. Schmidt, B. Brüne, and A. von Knethen, "The nuclear hormone receptor PPAR γ as a therapeutic target in major diseases," *The Scientific World Journal*, vol. 10, pp. 2181–2197, 2010.
- [76] J. M. Kaplan and B. Zingarelli, "Novel therapeutic agents in pediatric sepsis: peroxisome proliferator receptor γ (PPAR γ) agonists," *The Open Inflammation Journal*, vol. 4, no. 1, pp. 120–124, 2011.
- [77] A. C. C. Wang, X. Dai, B. Luu, and D. J. Conrad, "Peroxisome proliferator-activated receptor- γ regulates airway epithelial cell activation," *American Journal of Respiratory Cell and Molecular Biology*, vol. 24, no. 6, pp. 688–693, 2001.
- [78] K. S. Lee, S. J. Park, P. H. Hwang et al., "PPAR-gamma modulates allergic inflammation through up-regulation of PTEN," *The FASEB Journal*, vol. 19, no. 8, pp. 1033–1035, 2005.
- [79] S. R. Kim, K. S. Lee, H. S. Park et al., "Involvement of IL-10 in peroxisome proliferator-activated receptor gamma-mediated anti-inflammatory response in asthma," *MolPharmacol*, vol. 68, pp. 1568–1575, 2005.
- [80] K. Honda, P. Marquillies, M. Capron, and D. Dombrowicz, "Peroxisome proliferator-activated receptor γ is expressed in airways and inhibits features of airway remodeling in a mouse asthma model," *The Journal of Allergy and Clinical Immunology*, vol. 113, no. 5, pp. 882–888, 2004.
- [81] L. Benayoun, S. Letuve, A. Druilhe et al., "Regulation of peroxisome proliferator-activated receptor γ expression in human asthmatic airways: Relationship with proliferation, apoptosis, and airway remodeling," *American Journal of Respiratory and Critical Care Medicine*, vol. 164, no. 8 I, pp. 1487–1494, 2001.
- [82] M. Kobayashi, M. J. Thomassen, T. Rambasek et al., "An inverse relationship between peroxisome proliferator-activated receptor γ and allergic airway inflammation in an allergen challenge model," *Annals of Allergy, Asthma & Immunology*, vol. 95, no. 5, pp. 468–473, 2005.
- [83] J. E. Ward, D. J. Fernandes, C. C. Taylor, J. V. Bonacci, L. Quan, and A. G. Stewart, "The PPAR γ ligand, rosiglitazone, reduces airways hyperresponsiveness in a murine model of allergen-induced inflammation," *Pulmonary Pharmacology and Therapeutics*, vol. 19, no. 1, pp. 39–46, 2006.
- [84] V. R. Narala, R. Ranga, M. R. Smith et al., "Pioglitazone is as effective as dexamethasone in a cockroach allergen-induced murine model of asthma," *Respiratory Research*, vol. 8, article no. 90, 2007.
- [85] C. Mueller, V. Weaver, J. P. Vanden Heuvel, A. August, and M. T. Cantorna, "Peroxisome proliferator-activated receptor gamma ligands attenuate immunological symptoms of experimental allergic asthma," *Archives of Biochemistry and Biophysics*, vol. 418, no. 2, pp. 186–196, 2003.
- [86] S. Ueki, Y. Matsuwaki, H. Kayaba et al., "Peroxisome proliferator-activated receptor γ regulates eosinophil functions: A new therapeutic target for allergic airway inflammation," *International Archives of Allergy and Immunology*, vol. 134, no. 1, pp. 30–36, 2004.
- [87] V. Angeli, H. Hammad, B. Staels, M. Capron, B. N. Lambrecht, and F. Trottein, "Peroxisome proliferator-activated receptor γ inhibits the migration of dendritic cells: consequences for the immune response," *The Journal of Immunology*, vol. 170, no. 10, pp. 5295–5301, 2003.
- [88] H. Hammad, H. J. De Heer, T. Soullié et al., "Activation of Peroxisome Proliferator-Activated Receptor- γ in Dendritic Cells Inhibits the Development of Eosinophilic Airway Inflammation in a Mouse Model of Asthma," *The American Journal of Pathology*, vol. 164, no. 1, pp. 263–271, 2004.
- [89] K. Asada, S. Sasaki, T. Suda, K. Chida, and H. Nakamura, "Antiinflammatory roles of peroxisome proliferator-activated receptor gamma in human alveolar macrophages," *American Journal of Respiratory and Critical Care Medicine*, vol. 169, no. 2, pp. 195–200, 2004.
- [90] Y.-S. Yoon, S.-Y. Kim, M.-J. Kim, J.-H. Lim, M.-S. Cho, and J. L. Kang, "PPAR γ activation following apoptotic cell instillation promotes resolution of lung inflammation and fibrosis via regulation of efferocytosis and proresolving cytokines," *Mucosal Immunology*, vol. 8, no. 5, pp. 1031–1046, 2015.
- [91] H. J. Patel, M. G. Belvisi, D. Bishop-Bailey, M. H. Yacoub, and J. A. Mitchell, "Activation of peroxisome proliferator-activated receptors in human airway smooth muscle cells has a superior anti-inflammatory profile to corticosteroids: Relevance for chronic obstructive pulmonary disease therapy," *The Journal of Immunology*, vol. 170, no. 5, pp. 2663–2669, 2003.
- [92] M. Hetzel, D. Walcher, M. Grüb, H. Bach, V. Hombach, and N. Marx, "Inhibition of MMP-9 expression by PPAR γ activators in human bronchial epithelial cells," *Thorax*, vol. 58, no. 9, pp. 778–783, 2003.
- [93] K. S. Lee, S. J. Park, S. R. Kim et al., "Modulation of airway remodeling and airway inflammation by peroxisome proliferator-activated receptor γ in a murine model of toluene diisocyanate-induced asthma," *The Journal of Immunology*, vol. 177, no. 8, pp. 5248–5257, 2006.

- [94] T. J. Standiford, V. C. Keshamouni, and R. C. Reddy, "Peroxisome proliferator-activated receptor- γ as a regulator of lung inflammation and repair," *Proceedings of the American Thoracic Society*, vol. 2, no. 3, pp. 226–231, 2005.
- [95] J. E. Milam, V. G. Keshamouni, S. H. Phan et al., "PPAR- γ agonists inhibit profibrotic phenotypes in human lung fibroblasts and bleomycin-induced pulmonary fibrosis," *American Journal of Physiology-Lung Cellular and Molecular Physiology*, vol. 294, no. 5, pp. L891–L901, 2008.
- [96] H. A. Burgess, L. E. Daugherty, T. H. Thatcher et al., "PPAR γ agonists inhibit TGF- β induced pulmonary myofibroblast differentiation and collagen production: implications for therapy of lung fibrosis," *American Journal of Physiology-Lung Cellular and Molecular Physiology*, vol. 288, no. 6, pp. L1146–L1153, 2005.
- [97] S. T. Rinne, L. C. Feemster, B. F. Collins et al., "Thiazolidinediones and the risk of asthma exacerbation among patients with diabetes: a cohort study," *Allergy, Asthma & Clinical Immunology*, vol. 10, no. 1, p. 34, 2014.
- [98] M. S. Sandhu, V. Dimov, A. K. Sandhu, R. W. Walters, T. Wichman, and T. Casale, "The use of the peroxisome proliferator-activated receptors γ agonist rosiglitazone to treat airway hyperreactivity," *Annals of Allergy, Asthma & Immunology*, vol. 109, no. 1, pp. 75–77, 2012.
- [99] Y. Hashimoto and K. Nakahara, "Improvement of asthma after administration of pioglitazone.," *Diabetes Care*, vol. 25, no. 2, p. 401, 2002.
- [100] M. Spears, I. Donnelly, L. Jolly et al., "Bronchodilatory effect of the PPAR- γ agonist rosiglitazone in smokers with asthma," *Clinical Pharmacology & Therapeutics*, vol. 86, no. 1, pp. 49–53, 2009.
- [101] D. B. Richards, P. Bareille, E. L. Lindo, D. Quinn, and S. N. Farrow, "Treatment with a peroxisomal proliferator activated receptor gamma agonist has a modest effect in the allergen challenge model in asthma: A randomised controlled trial," *Respiratory Medicine*, vol. 104, no. 5, pp. 668–674, 2010.
- [102] J. R. Anderson, K. Mortimer, L. Pang et al., "Evaluation of the PPAR- γ agonist pioglitazone in mild asthma: A double-blind randomized controlled trial," *PLoS ONE*, vol. 11, no. 8, Article ID e0160257, 2016.
- [103] B. D. Levy, P. J. Noel, M. M. Freemer et al., "Future research directions in asthma: An NHLBI working group report," *American Journal of Respiratory and Critical Care Medicine*, vol. 192, no. 11, pp. 1366–1372, 2015.
- [104] R. W. Nesto, D. Bell, R. O. Bonow et al., "Thiazolidinedione use, fluid retention, and congestive heart failure: a consensus statement from the American Heart Association and American Diabetes Association," *Diabetes Care*, vol. 27, no. 1, pp. 256–263, 2004.
- [105] D. J. Betteridge, "Thiazolidinediones and fracture risk in patients with Type 2 diabetes," *Diabetic Medicine*, vol. 28, no. 7, pp. 759–771, 2011.
- [106] A. J. Scheen, "Thiazolidinediones and liver toxicity," *Diabetes & Metabolism*, vol. 27, no. 3, pp. 305–313, 2001.
- [107] A. Ialenti, G. Grassia, P. Di Meglio, P. Maffia, M. Di Rosa, and A. Ianaro, "Mechanism of the anti-inflammatory effect of thiazolidinediones: Relationship with the glucocorticoid pathway," *Molecular Pharmacology*, vol. 67, no. 5, pp. 1620–1628, 2005.
- [108] S. Fogli, F. Stefanelli, L. Picchianti et al., "Synergistic interaction between PPAR ligands and salbutamol on human bronchial smooth muscle cell proliferation," *British Journal of Pharmacology*, vol. 168, no. 1, pp. 266–275, 2013.
- [109] M. Nie, L. Corbett, A. J. Knox, and L. Pang, "Differential regulation of chemokine expression by peroxisome proliferator-activated receptor γ agonists: Interactions with glucocorticoids and β 2-agonists," *The Journal of Biological Chemistry*, vol. 280, no. 4, pp. 2550–2561, 2005.